

Distance Metrics.*

Greg Kochanski

<http://kochanski.org/gpk>

July 25, 2009

To talk about “close” and “far” in a consistent manner, you need to be able to measure distances. For this, you may need the concept of a “distance metric.”

A distance metric is a little bit more than something that starts at zero and gets bigger as you get farther away. It must be defined in such a way that the shortest distance between any two points is a straight line.

A distance metric, $d(A, B)$ is a function or algorithm for calculating a distance between two things, A and B . It has three properties:

1. It is always positive or zero.
2. The distance from a thing¹ to itself is zero.
3. It obeys the triangle inequality: For any three points, A , B , and C , $d(A, B) + d(B, C) \geq d(A, C)$ for any possible choice of B . In other words, the straight line between A and C , which has a length $d(A, C)$, is shorter² than any other path between A and C , such as a path that goes by way of B .

Anything that obeys these three properties is a distance metric.

The standard examples are Euclidean distance in two dimensions:

$$d_{Euclid}(A, B) = ((A_x - B_x)^2 + (A_y - B_y)^2)^{1/2}, \quad (1)$$

and the two-dimensional city block metric [Krowne, 2003]:

$$d_{Euclid}(A, B) = |A_x - B_x| + |A_y - B_y|. \quad (2)$$

Both Euclidean and city-block metrics can work in more (or less) than two dimensions. In either case, you get one term in the addition for each dimension.

References

[Krowne, 2003] Krowne, A. (2003). *PlanetMath: city-block metric*. PlanetMath.org, <http://planetmath.org/encyclopedia/CityBlockMetric.html>. Canonical name CityBlockMetric,.

*This work is licensed under the Creative Commons Attribution License. To view a copy of this license, visit <http://creativecommons.org/licenses/by/1.0/> or send a letter to Creative Commons, 559 Nathan Abbott Way, Stanford, California 94305, USA. This work is available under <http://kochanski.org/gpk/teaching/0401Oxford>.

¹ A geometrical point, or a document....

² Strictly speaking this should read “. . . at least as short as. . .,” because if B is on the line between A and C , then $d(A, B) + d(B, C) = d(A, C)$. Also, it’s possible to have a distance measure where certain directions just don’t count: an extreme example is if $d(X, Y) = 0$ for all X and Y . This always-zero (and rather useless) distance metric obeys the triangle inequality because $d(A, B) + d(B, C) = d(A, C)$ for all B .